



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2020

---

## Dealing with DNA lesions: When one cell cycle is not enough

Lezaja, Aleksandra ; Altmeyer, Matthias

**Abstract:** Subversion of genome integrity fuels cellular adaptation and is a prerequisite for organismal evolution, yet genomic lesions are also the harmful driving force of cancer and other age-related human diseases. Genome integrity maintenance is inherently linked to genome organization and nuclear architecture, which are substantially remodeled during the cell cycle. Here we discuss recent findings on how actively dividing cells cope with endogenous genomic lesions that occur frequently at repetitive, heterochromatic, and late replicating regions as byproducts of genome duplication. We discuss how such lesions, rather than being resolved immediately when they occur, are dealt with in subsequent cell cycle phases, and even after mitotic cell division, and how this in turn affects genome organization, stability, and function.

DOI: <https://doi.org/10.1016/j.ceb.2020.11.001>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-201550>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Lezaja, Aleksandra; Altmeyer, Matthias (2020). Dealing with DNA lesions: When one cell cycle is not enough. *Current Opinion in Cell Biology*, 70:27-36.

DOI: <https://doi.org/10.1016/j.ceb.2020.11.001>



# Dealing with DNA lesions: When one cell cycle is not enough

Aleksandra Lezaja and Matthias Altmeyer

## Abstract

Subversion of genome integrity fuels cellular adaptation and is a prerequisite for organismal evolution, yet genomic lesions are also the harmful driving force of cancer and other age-related human diseases. Genome integrity maintenance is inherently linked to genome organization and nuclear architecture, which are substantially remodeled during the cell cycle. Here we discuss recent findings on how actively dividing cells cope with endogenous genomic lesions that occur frequently at repetitive, heterochromatic, and late replicating regions as byproducts of genome duplication. We discuss how such lesions, rather than being resolved immediately when they occur, are dealt with in subsequent cell cycle phases, and even after mitotic cell division, and how this in turn affects genome organization, stability, and function.

## Addresses

Department of Molecular Mechanisms of Disease, University of Zurich, Zurich, Switzerland

Corresponding author: Altmeyer, Matthias ([matthias.altmeyer@uzh.ch](mailto:matthias.altmeyer@uzh.ch))

**Current Opinion in Cell Biology** 2021, 70:27–36

This review comes from a themed issue on **Cell Nucleus**

Edited by **Jane Skok** and **Daniel Gerlich**

For a complete overview see the [Issue](#) and the [Editorial](#)

<https://doi.org/10.1016/j.ceb.2020.11.001>

0955-0674/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Keywords

Genome stability, Cell cycle control, Mitosis, MiDAS, Telomere maintenance, ALT, Chromatin compartments, Liquid–liquid phase separation.

## Introduction

Among the main tasks of the cell nucleus are the functional organization and stabilization of the genome. This involves the spatial organization of DNA into nucleosomes, nucleosome clusters, chromatin loops, topologically associating domains (TADs), larger chromosome compartments, and chromosome territories [1,2]. But it also involves the constant remodeling of genome architecture to facilitate genome functions such as gene

expression, DNA replication, cell division, and DNA repair. In this sense, the term *genome architecture* is misleading, as the genome is far from being static. Indeed, when genome organization is analyzed at single cell resolution, a continuous cell cycle–dependent reorganization of chromatin domains is revealed [3]. Genome duplication during S-phase of the cell cycle, chromosome condensation during mitotic cell division, and chromosome decondensation in early G1 all represent major, yet in their spatiotemporal regulation incompletely understood, genome remodeling events. Maintaining genome integrity throughout these processes is critical for cell function, and failure to do so is associated with accumulation of mutations, chromosomal aberrations, and cellular transformation leading to disease.

Genome integrity maintenance is the primary function of two interconnected cellular networks, the DNA damage response (DDR) [4,5] and the replication stress response (RSR) [6]. While the DDR takes care of chromosome breaks and other types of DNA lesions throughout the cell cycle, the RSR is primarily active in S-phase to deal with stalled replication forks and harmful replication intermediates. Although they are typically considered distinct cellular responses and often studied separately, emerging evidence suggests that certain replication intermediates, such as reversed replication forks, structurally resemble DNA lesions and that the DDR and the RSR employ a shared set of molecular players to maintain a stable genome [7]. Furthermore, both the DDR and the RSR trigger cell cycle checkpoints to slow down or halt cell cycle progression, and tight connections exist between the mechanisms involved in genome integrity maintenance and cell cycle control [8,9]. On the other hand, cell cycle checkpoints are not failsafe. Accumulating evidence suggests that replication intermediates and genomic lesions originating from endogenous replication stress can escape checkpoint surveillance and are transmitted to subsequent cell cycle phases and even to the next cell generation [10,11]. In this review we highlight and discuss regions in the genome that are particularly vulnerable to endogenous replication stress and thus inherently fragile, and provide an update on the emerging mechanisms used by cells to deal with this fragility in a manner that encompasses multiple successive cell cycle phases. As the genome undergoes

significant reorganization when cells cross the borders between cell cycle phases, for example, when they progress from S/G2 into mitosis and from mitosis into G1, we will put these mechanisms into the context of nuclear compartmentalization and discuss how this in turn may impact genome stability. Although dealing with DNA lesions across cell cycle boundaries is not limited to cancer cells, it may be particularly important as a source of cancer driving mutations, to shape cancer genome evolution, and to link deregulated genome integrity maintenance and genotoxic cancer therapies to innate immunity. Understanding the underlying principles and to which extent cancer cells rely on them may therefore offer new opportunities for therapeutic interventions.

### Endogenous replication stress and inherently fragile regions in the genome


Replication stress arises when replication fork progression is impaired [6]. This can occur in response to DNA lesions caused by exogenous DNA damaging agents, including many chemotherapeutic drugs, but it can also occur during normal S-phase progression in the absence of exogenous stress. Frequent replication fork slowing and transient pausing thus appear to be natural byproducts of genome duplication [7]. Among the sources for such endogenous replication stress are conflicts between the replication and transcription

machineries, unusual DNA structures such as G-quadruplexes, bulky DNA-protein complexes, and difficult-to-replicate regions in the genome. The latter include chromosomal fragile regions, so-called common and rare fragile sites, as well as repetitive sequences at telomeres, centromeres, and ribosomal DNA (rDNA) (Figure 1). In addition to being inherently difficult to replicate, these regions are also replicated late in the cell cycle, and, as we will discuss in more detail below, their late replication timing significantly shortens the time window available in S-phase to complete their replication. This in turn primes them for replication carried on beyond S-phase, by mechanisms, which on one hand provide a safety net for completion of genome duplication but which on the other hand also pose a threat to genome integrity and render these regions particularly vulnerable.

### Chromosomal fragile sites

Chromosomal fragile sites are hotspots for DNA breakage and aberrations such as deletions, duplications, and translocations in response to replication stress [12]. Based on their prevalence, fragile sites are typically classified as either rare or common (Figure 1). While rare fragile sites, such as the CGG triplet expansion in the *FMR1* gene underlying fragile X syndrome, are found only in a small percentage of the

Figure 1



**Fragile genomic regions.** Repetitive, heterochromatic, and late replicating regions in the genome are inherently fragile and frequently experience replication stress-associated damage. Fragile genomic regions include telomeres at chromosome ends and subtelomeric repeats, centromeres and pericentromeric repeats, rDNA repeats in the nucleolus, as well as common and rare fragile sites.

population, common fragile sites (CFSs), such as *FRA16D* and *FRA3B* with the associated tumor suppressor genes *WWOX* and *FHIT*, respectively, are present in the general population and frequently mutated in cancer [13]. CFSs are typically late replicating, origin-scarce regions. They are composed of AT-rich sequences, which can form secondary structures that stall replication and impair repair [14], and they are associated with very long genes at TAD boundaries that can take more than one cell cycle to be fully transcribed, inevitably evoking transcription-replication conflicts [15,16]. This combination of challenges impedes faithful and complete CFSs replication during S-phase, leading to under-replicated DNA and endangered CFSs stability [17]. Of note, transcription and replication timing are intertwined, and it was recently demonstrated that experimentally enhanced transcription, induced by a strong promoter, triggers a switch to earlier replication timing and alleviates fragility [18]. While this reinforces the notion that late replication of CFSs underlies their fragility, a distinct class of early replicating fragile sites in highly expressed gene clusters has also been identified [19,20••]. Their fragility is consistent with the recent finding that oncogene activation induces firing of normally suppressed replication origins within highly expressed genes, causing transcription-replication interference upon premature S-phase entry [21••]. Fragility thus not only occurs when time gets short at the end of S-phase but also at the transition from G1 to S-phase when high transcriptional activity collides with the rapid firing of origins to initiate replication.

### Telomeres

Telomeres and the adjacent subtelomeric repeats are the regions that protect chromosome ends (Figure 1). In human cells, telomeres are composed of terminal TTAGGG repeat sequences, which can span several kilobases in length and which are characterized by constitutive heterochromatin. Despite their heterochromatic nature, the long noncoding RNA TERRA is transcribed from subtelomeric repeats and forms telomere-associated R-loops that can interfere with telomere replication and stability [22,23]. Secondary structures such as G-quadruplexes formed at telomeric G-rich repetitive sequences also challenge replication fork progression [24,25]. Replication stress at telomeres is further enhanced due to unidirectional replication downstream of the most distal origin, which means that irreversibly stalled replication forks cannot be rescued by dormant origin firing. Finally, the telomere-associated t-loop formed by the terminating single-stranded G-rich overhang and the tightly bound proteins of the protective shelterin complex challenge replisome progression and make telomere replication inherently difficult [26].

Telomere length maintenance, which is needed to overcome the end replication problem and achieve replicative immortality, depends on the reverse transcriptase telomerase, or in around 10–15% of cancers (including gliomas, sarcomas, and pancreatic neuroendocrine tumors), on alternative lengthening of telomeres (ALT) [24,27]. Telomere elongation by ALT relies on break-induced replication, a recombination-driven process that is initiated in the context of DNA replication but then continues as break-induced telomere synthesis beyond S-phase [28–33]. Break-induced replication additionally contributes to telomere fragility, and cancer cells using ALT for telomere maintenance show higher levels of telomere-associated replication stress compared to telomerase expressing non-ALT cancer cells [22,24,34••,35•].


### Centromeres

Centromeres are the assembly sites for the kinetochore, the protein complex that makes the connection between chromosomes and the spindle fibers, so that the chromatids can be pulled toward opposite poles during cell division (Figure 1). In human cells, centromeres are composed of a series of AT-rich head-to-tail tandem repeats named *alpha satellites*, which extend for several megabases [36]. Furthermore, the surrounding pericentromeric DNA is also organized in short tandem repeats. The repetitive nature of centromeric and pericentromeric sequences, the formation of DNA secondary structures, the heterochromatin environment and their generally late replication timing all contribute to centromere fragility [37]. Due to their central role for chromosome segregation, centromeric and pericentromeric replication stress, chromosome breaks and recombination are associated with chromosomal aberrations and translocations in cancer.

### rDNA

Ribosomal DNA contains several hundred copies of repeated gene sequences (Figure 1). The ribosomal rRNAs, required for ribosome biosynthesis, are transcribed from these rDNA repeats, making them the most heavily transcribed regions in the genome. The human rDNA repeats encompass several genomic loci, with the 5 S rRNA gene repeats encoded by chromosome 1, and the 18 S, 5.8 S, and 28 S rRNA gene repeats distributed over chromosomes 13–15 and 21–22 [38]. These head-to-tail repeat clusters on multiple chromosomes come together in the nucleolus, a membraneless nuclear compartment. Due to an excess number of rDNA repeats, not all repeats are transcribed into rRNA. Silent rDNA repeats form tightly packed heterochromatin at the nucleolar periphery and, unlike actively transcribed repeats within the nucleolus, are replicated

Figure 2



**Dealing with DNA lesions across cell cycle boundaries.** While DNA replication is normally confined to the S-phase of the cell cycle, in response to endogenous and exogenous replication stress, e.g. upon oncogene activation or loss of tumor suppressor genes, replication and recombination intermediates are transmitted to later stages of the cell cycle. Late replicating regions, in which replication problems occur late in S-phase, are predestined for continuing replication in S/G2, and at under-replicated DNA replication continues even in mitosis. Mitotic DNA synthesis (MiDAS) is a conservative form of break-induced replication that works at common fragile sites and telomeres. Upon exit from mitosis, unresolved replication intermediates and chromosome breaks can give rise to micronuclei and free cytoplasmic DNA, such as extrachromosomal telomeric repeat DNA and C-circles, which in turn can activate the cGAS-STING pathway and trigger a type I interferon response. In the context of cancer treatment, chemo- and radiation therapy (including genotoxic cell cycle and DNA repair drugs such as ATR, WEE1, and PARP inhibitors) may thus synergize with immune response activating checkpoint inhibitors to maximize cancer cell death. In the newly forming main nucleus, 53BP1 condensates shield inherited DNA damage and are associated with clustering of difficult-to-repair lesions and a p53-p21-response. This extends G1 duration in response to replication stress during the previous cell cycle and determines whether and when cells commit to the next round of replication. In absence of such failsafe mechanisms (e.g. upon p53 or p21 loss), transgenerational lesion repair likely leads to a progressive accumulation of mutations and fuels cancer evolution.

late in S-phase. Rearrangements of rDNA repeats, rDNA repeat expansions and contractions, and rDNA-associated structural chromosomal aberrations are frequently found in cancer. The instability of rDNA in disease is likely driven by transcription-replication conflicts due to R-loops at transcribed rDNA repeats, and by replication impediments coming from silent, heterochromatic rDNA repeats and from their late replication timing.

### Dealing with incompletely replicated DNA beyond S-phase

As mentioned above, inherently fragile regions may not only be susceptible to replication stress and DNA damage because of their difficult-to-replicate sequence but also because of their late replication timing. In particular in cancer cells, where cell cycle checkpoint control is loosened and premature cell cycle transitions occur, late replicating regions may continue replication



in G2 and even mitosis (Figure 2). Indeed, G2 could be viewed as an extended S-phase [39], as a time in the cell cycle when late replicating and under-replicated DNA is taken care of. Replication stress, in particular at origin scarce regions where fork stalling cannot be resolved by dormant origin firing, exacerbates the amount of incompletely replicated DNA. Likewise, shortening the S/G2 phase by overriding intrinsic checkpoint control and enforcing premature activation of mitotic processes, e.g. through inhibition of the ATR, CHK1, and WEE1 kinases [40–45], increases the fraction of the genome that has not been replicated fully. How do cells deal with such under-replicated DNA? Surprisingly, DNA replication does not seem to stop upon mitotic entry, as accumulating evidence has shown [46–52]. Mitotic DNA synthesis (MiDAS) occurs at CFSs and fragile telomeres and uses a form of break-induced replication. This conservative form of DNA synthesis depends on the replicative polymerase  $\delta$  and its subunit POLD3, on the scaffolding factor SLX4, the MUS81-EME1 endonuclease, and on the RAD52 recombinase [53–58]. The switch from semi-conservative replication in S/G2 to MiDAS also seems to require prior replisome disassembly, which is mediated by the TRAIP ubiquitin ligase and CDC48/p97-dependent extraction of the replicative helicase from chromatin [59–62]. While the mechanism of MiDAS at CFSs and telomeres and the main players involved have started to emerge, to which extent MiDAS occurs at centromeres, rDNA repeats, and other fragile regions, and how different subpathways of MiDAS may deal with different fragile loci are currently not very well understood.

As genome duplication and cell division should be temporally separated in order to ensure that the complete genetic information is transmitted to the offspring, MiDAS may be the last resort to complete replication before chromosome segregation. Challenging this paradigm, it was recently shown in budding yeast that DNA synthesis is inhibited in metaphase due to high CDK activity, and allowed to resume during anaphase, late in mitosis when chromosomes separate [63•]. As mammalian MiDAS seems restricted to the time window from prophase to metaphase, whether post-metaphase DNA synthesis exists in human cells is currently not known.

What happens when MiDAS fails, or time is simply too short during mitosis to complete replication of under-replicated DNA coming from S/G2? Unresolved replication intermediates can lead to mitotic chromatin bridges, lagging chromosomes, nucleosome-free ultra-fine anaphase bridges, and chromosome breaks [64–66]. These aberrations can give rise to 53BP1 nuclear bodies in G1 cells, which shield inherited genomic lesions from nucleolytic degradation and unscheduled repair reactions, and they can lead to broken DNA escaping from the newly forming nucleus to generate micronuclei and

extrachromosomal DNA (Figure 2). Recent work has started to shed light on the consequences of such inherited genomic lesions, linking genome integrity maintenance to nuclear organization by liquid condensates and to cGAS-STING-mediated activation of innate immunity.

### Consequences of replication stress-associated heritable DNA lesions

The time from nuclear envelope breakdown in prophase to its reformation in telophase poses a particular threat to genome integrity maintenance not only because of the massive chromatin rearrangements that happen during mitosis but also because the physical barrier that normally prevents leakage of DNA into the cytoplasm is gone. Although sophisticated mechanisms are in place to keep mitotic chromosomes spatially constrained and to separate nuclear from cytoplasmic components during nuclear reassembly [67•,68], broken pieces of chromosomes can escape from the newly forming nucleus to constitute micronuclei and cytoplasmic extrachromosomal DNA (Figure 2). Micronuclei have a fragile envelope that easily breaks down during interphase and they show asynchronous replication, which on one hand can lead to chromosome aberrations such as the complex rearrangements seen in chromothripsis, and on the other hand trigger, through cytoplasmic self-DNA sensing by the cGAS-STING pathway, an innate immune response [69•,70,71]. Telomere fragility and dysfunction during replicative crisis as well as extrachromosomal telomeric repeat DNA have also been associated with chromothripsis and cGAS-STING activation [72,73•,74•]. Nuclear escape of DNA late in mitosis as a consequence of replication stress and incomplete MiDAS at fragile genomic regions could therefore be a general mechanism that links deregulated or impaired genome integrity maintenance to genomic rearrangements and innate immune signaling (Figure 2).

The cGAS-STING pathway is also activated in cells that lost the tumor suppressors and homology-directed repair factors BRCA1 or BRCA2 [75–79]. Importantly, chemotherapeutic drugs that exacerbate replication stress and DNA damage, such as PARP and ATR inhibitors, result in more problems late in the cell cycle and during mitosis [41•,80•], and further stimulate the cGAS-STING-mediated type I interferon response, suggesting that such compounds can be exploited in certain contexts to boost cancer immunotherapies [81,82].

While mitotic exit bears the risk of losing damaged DNA from the newly forming nucleus, re-establishing the nuclear compartment also provides new opportunities to deal with inherited DNA lesions on correctly segregated chromosomes. The multivalent chromatin reader 53BP1 simultaneously senses replication status and presence of

DNA damage and through its oligomerization domain forms large compartments around genomic lesions in G1 cells (Figure 2). These protein condensates, which promote repair and stabilize the chromatin topology in the vicinity of DNA lesions [83••], show many features of liquid–liquid phase separation (LLPS), including concentration-dependent self-assembly and condensate fusions [84••,85••]. They are precluded from forming during mitosis, when the 53BP1 concentration is diluted after nuclear envelope breakdown and when high kinase activities, including CDK-dependent phosphorylations, prevent LLPS and 53BP1 assembly [86,87]. In G1, however, 53BP1 avidly assembles at inherited lesions and, via p53 and its downstream target p21, modulates G1 length and S-phase commitment [88–90]. Difficult-to-repair lesions inside such condensates are particularly prone to undergo clustering by compartment fusions [84••,91,92•], resulting in repair centers that facilitate repair but also bear the risk of promoting chromosomal translocations. The role of 53BP1 extends beyond G1, as it also impacts replication fork stability, replication timing of inherited lesions, and coordinates the hand-over to RAD52-mediated recombination later in the cell cycle [93•,94•].

While 53BP1 condensates are primarily associated with under-replicated DNA at CFSs, LLPS-mediated clustering has recently been described for inherently fragile telomeres in ALT-positive cancer cells [95•,96•]. ALT telomeres, mediated by the SUMO pathway, assemble into ALT-associated PML bodies (APBs), liquid-like condensates in which inter- and intrachromosomal break-induced replication occurs. As PML bodies are also recruited to persistent ionizing radiation-induced DNA lesions [97], compartmentalization into nuclear condensates may emerge as a common theme for a broader range of inherently fragile as well as difficult-to-repair genomic regions.

## Conclusions

As opposed to the classic cell cycle arrest, repair, and restart model, work from recent years has started to support the notion that certain genomic lesions take more than one cell cycle phase and sometimes even more than one cell cycle to be repaired. Particularly replication stress-associated lesions and physiological replication intermediates, such as stretches of under-replicated DNA at late replicating regions in the genome, have a comparatively short time-window for repair in S-phase, and they depend on replication and repair events in late S/G2 and mitosis. This makes them especially fragile with consequences for genome stability and for the detection of DNA damage by the immune system.

Mutational cascades amplifying chromothripsis over successive cell cycles, and segregation of unrepaired

genomic lesions over multiple cell generations drives evolution, subclonal heterogeneity, and adaptation of cancer genomes [98••,99••]. Furthermore, nonrandom segregation of DNA lesions in response to replication stress was recently shown to cause biased inheritance of newly replicated DNA by daughter cells [100••]. A better understanding of the underlying mechanisms and of the dynamic evolution of replication stress phenotypes may not only yield information on the origin and evolution of cancer but also hold promises for more targeted treatments. ATR inhibitors, for instance, enhance asymmetric distribution of replication stress-associated DNA damage [100••], and potentiate cGAS-STING-mediated interferon signaling [81]. In targeted cancer therapy, these two effects might synergize for desired outcomes.

While CFSs were the first fragile regions to be linked to 53BP1 nuclear bodies and MiDAS, telomeric MiDAS has recently received more attention. However, it is not entirely clear yet how ALT-negative and ALT-positive cancer cells differ in terms of MiDAS and replication stress-associated telomere fragility. The fragility of other genomic regions, including centromeres, pericentromeric repeats, and rDNA arrays, is even less well understood. New technologies that can map fragile regions and their replication genome-wide in specific cell cycle phases have recently been developed [101•,102•,103•], and they will likely shed new light on the mechanism of MiDAS-dependent recombination repair and also identify additional fragile regions. Applied to different cell types with different sets of activated oncogenes and inactivated tumor suppressors, a more comprehensive and condition-specific view of the fragile genome will be obtained. Complemented by time-resolved single cell experiments to investigate the fate of individual inherited DNA lesions for the next cell generations, we will understand better how cells deal with DNA damage when one cell cycle is not enough.

## Conflict of interest statement

Nothing declared.

## Acknowledgements

We apologize to all colleagues whose work could not be cited due to space limitations, and acknowledge research funding from the Swiss National Science Foundation (PP00P3\_179057), the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (ERC-2016-STG 714326), the Novartis Foundation for Medical-Biological Research (16B078), and the Swiss Foundation to Combat Cancer (Stiftung zur Krebsbekämpfung). The authors declare no conflict of interest.

## References

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Szabo Q, Bantignies F, Cavalli G: **Principles of genome folding into topologically associating domains**. *Science Advances* 2019, **5**.
2. Yu M, Ren B: **The three-dimensional organization of mammalian genomes**. *Annu Rev Cell Dev Biol* 2017, **33**: 265–289.
3. Nagano T, Lubling Y, Vaarnai C, Dudley C, Leung W, Baran Y, Cohen NM, Wingett S, Fraser P, Tanay A: **Cell-cycle dynamics of chromosomal organization at single-cell resolution**. *Nature* 2017, **547**:61–67.
4. Jackson SP, Bartek J: **The DNA-damage response in human biology and disease**. *Nature* 2009, **461**:1071–1078.
5. Ciccia A, Elledge SJ: **The DNA damage response: making it safe to play with knives**. *Mol Cell* 2010, **40**:179–204.
6. Zeman MK, Cimprich KA: **Causes and consequences of replication stress**. *Nat Cell Biol* 2014, **16**:2–9.
7. Berti M, Cortez D, Lopes M: **The plasticity of DNA replication forks in response to clinically relevant genotoxic stress**. *Nat Rev Mol Cell Biol* 2020, **21**:633–651.
8. Murray JM, Carr AM: **Integrating DNA damage repair with the cell cycle**. *Curr Opin Cell Biol* 2018, **52**:120–125.
9. Hustedt N, Durocher D: **The control of DNA repair by the cell cycle**. *Nat Cell Biol* 2016, **19**:1–9.
10. Mankouri HW, Huttner D, Hickson ID: **How unfinished business from S-phase affects mitosis and beyond**. *EMBO J* 2013, **32**: 2661–2671.
11. Lukas J, Lukas C, Bartek J: **More than just a focus: the chromatin response to DNA damage and its role in genome integrity maintenance**. *Nat Cell Biol* 2011, **13**:1161–1169.
12. Irony-Tur Sinai M, Kerem B: **Genomic instability in fragile sites—still adding the pieces**. *Genes Chromosomes Cancer* 2019, **58**: 295–304.
13. Ozer O, Hickson ID: **Pathways for maintenance of telomeres and common fragile sites during DNA replication stress**. *Open Biology* 2018, **8**.
14. Kaushal S, Wollmuth CE, Das K, Hile SE, Regan SB, Barnes RP, Haouzi A, Lee SM, House NCM, Guyumdzhyan M, et al.: **Sequence and nuclease requirements for breakage and healing of a structure-forming (AT)<sub>n</sub> sequence within fragile site FRA16D**. *Cell Rep* 2019, **27**:1151–1164.
15. Brison O, El-Hilali S, Azar D, Koundrioukoff S, Schmidt M, Nahse V, Jaszczyszyn Y, Lachages AM, Dutrillaux B, Thermes C, et al.: **Transcription-mediated organization of the replication initiation program across large genes sets common fragile sites genome-wide**. *Nat Commun* 2019, **10**.
16. Sami D, Sasaki T, Tur-Sinai MI, Miron K, Rivera-Mulia JC, Magnuson B, Ljungman M, Gilbert DM, Kerem B: **3D genome organization contributes to genome instability at fragile sites**. *Nat Commun* 2020, **11**.
17. Debatisse M, Rosselli F: **A journey with common fragile sites: from S phase to telophase**. *Genes Chromosomes Cancer* 2019, **58**:305–316.
18. Blin M, Le Tallec B, Nahse V, Schmidt M, Brossas C, Millot GA, Prioleau MN, Debatisse M: **Transcription-dependent regulation of replication dynamics modulates genome stability**. *Nat Struct Mol Biol* 2019, **26**:58–66.
19. Barlow JH, Faryabi RB, Callen E, Wong N, Malhowski A, Chen HT, Gutierrez-Cruz G, Sun HW, McKinnon P, Wright G, et al.: **Identification of early replicating fragile sites that contribute to genome instability**. *Cell* 2013, **152**:620–632.
20. Tubbs A, Sridharan S, van Wietmarschen N, Maman Y, Callen E, Stanlie A, Wu W, Wu X, Day A, Wong N, et al.: **Dual roles of poly(dA:dT) tracts in replication initiation and fork collapse**. *Cell* 2018, **174**:1127–1142. e1119.
- This study identifies poly(dA:dT) tracts as common feature of replication fork collapse at early replicating fragile sites, common fragile sites, and fragile rDNA repeats, and suggests that they promote replication initiation at the cost of chromosome fragility due to unprotected single-stranded poly(dA).
21. Macheret M, Halazonetis TD: **Intragenic origins due to short G1 phases underlie oncogene-induced DNA replication stress**. *Nature* 2018, **555**:112–116.
- This study, by analyzing nascent DNA synthesis in early S-phase, identifies thousands of new replication initiation sites upon oncogene activation that fall within highly transcribed genes, providing a mechanism for oncogene-induced replication stress due to transcription-replication interference.
22. Arora R, Lee Y, Wischniewski H, Brun CM, Schwarz T, Azzalin CM: **RNaseH1 regulates TERRA-telomeric DNA hybrids and telomere maintenance in ALT tumour cells**. *Nat Commun* 2014, **5**.
23. Azzalin CM, Reichenbach P, Khoraiu L, Giulotto E, Lingner J: **Telomeric repeat-containing RNA and RNA surveillance factors at mammalian chromosome ends**. *Science* 2007, **318**: 798–801.
24. Sobinoff AP, Pickett HA: **Alternative lengthening of telomeres: DNA repair pathways converge**. *Trends Genet* 2017, **33**: 921–932.
25. Vannier JB, Pavicic-Kaltenbrunner V, Petalcorin MI, Ding H, Boulton SJ: **RTEL1 dismantles T loops and counteracts telomeric G4-DNA to maintain telomere integrity**. *Cell* 2012, **149**: 795–806.
26. Sarek G, Kotsantis P, Ruis P, Van Ly D, Margalef P, Borel V, Zheng XF, Flynn HR, Snijders AP, Chowdhury D, et al.: **CDK phosphorylation of TRF2 controls t-loop dynamics during the cell cycle**. *Nature* 2019, **575**:523–527.
27. Hoang SM, O'Sullivan RJ: **Alternative lengthening of telomeres: building bridges to connect chromosome ends**. *Trends in Cancer* 2020, **6**:247–260.
28. Zhang JM, Yadav T, Ouyang J, Lan L, Zou L: **Alternative lengthening of telomeres through two distinct break-induced replication pathways**. *Cell Rep* 2019, **26**:955–968. e953.
29. Zhang T, Zhang Z, Shengzhao G, Li X, Liu H, Zhao Y: **Strand break-induced replication fork collapse leads to C-circles, C-overhangs and telomeric recombination**. *PLoS Genet* 2019, **15**, e1007925.
30. Dille RL, Verma P, Cho NW, Winters HD, Wondisford AR, Greenberg RA: **Break-induced telomere synthesis underlies alternative telomere maintenance**. *Nature* 2016, **539**:54–58.
31. Zhang JM, Zou L: **Alternative lengthening of telomeres: from molecular mechanisms to therapeutic outlooks**. *Cell Biosci* 2020, **10**:30.
32. Min J, Wright WE, Shay JW: **Alternative lengthening of telomeres mediated by mitotic DNA synthesis engages break-induced replication processes**. *Mol Cell Biol* 2017, **37**.
33. Roumelioti FM, Sotiropoulos SK, Katsini V, Chiourea M, Halazonetis TD, Gagos S: **Alternative lengthening of human telomeres is a conservative DNA replication process with features of break-induced replication**. *EMBO Rep* 2016, **17**: 1731–1737.
34. Kim J, Sturgill D, Sebastian R, Khurana S, Tran AD, Edwards GB, Kruswick A, Burkett S, Hosogane EK, Hannon WW, et al.: **Replication stress shapes a protective chromatin environment across fragile genomic regions**. *Mol Cell* 2018, **69**: 36–47. e37.
- This study demonstrates that the histone variant macroH2A1.2 is deposited at fragile genomic regions that experience recurrent replication stress. Over multiple cell generations, this generates macroH2A1.2 domains around fragile sites, which promote repair factor recruitment and protect against replicative senescence.
35. Kim J, Sun C, Tran AD, Chin PJ, Ruiz PD, Wang K, Gibbons RJ, Gamble MJ, Liu Y, Oberdoerffer P: **The macroH2A1.2 histone variant links ATRX loss to alternative telomere lengthening**. *Nat Struct Mol Biol* 2019, **26**:213–219.
- This study extends the protective role of the histone variant macroH2A1.2 to ALT telomeres, where macroH2A1.2 is dynamically



exchanged upon acute replication stress and facilitates homology-directed telomere repair.

36. Barra V, Fachinetti D: **The dark side of centromeres: types, causes and consequences of structural abnormalities implicating centromeric DNA.** *Nat Commun* 2018, **9**:4340.
  37. Mitrentsi I, Yilmaz D, Soutoglou E: **How to maintain the genome in nuclear space.** *Curr Opin Cell Biol* 2020, **64**:58–66.
  38. Warmerdam DO, Wolthuis RMF: **Keeping ribosomal DNA intact: a repeating challenge.** *Chromosome Res* 2019, **27**:57–72.
  39. Galarreta A, Lecona E, Valledor P, Ubieto P, Lafarga V, Specks J, Fernandez-Capetillo O: **USP7 couples DNA replication termination to mitotic entry.** *bioRxiv* 2018:305318, <https://doi.org/10.1101/305318>.
  40. Michelena J, Gatti M, Teloni F, Imhof R, Altmeyer M: **Basal CHK1 activity safeguards its stability to maintain intrinsic S-phase checkpoint functions.** *J Cell Biol* 2019, **218**:2865–2875.
  41. Michelena J, Lezaja A, Teloni F, Schmid T, Imhof R, Altmeyer M: **Analysis of PARP inhibitor toxicity by multidimensional fluorescence microscopy reveals mechanisms of sensitivity and resistance.** *Nat Commun* 2018, **9**.
  42. Lemmens B, Hegarat N, Akopyan K, Sala-Gaston J, Bartek J, Hochegger H, Lindqvist A: **DNA replication determines timing of mitosis by restricting CDK1 and PLK1 activation.** *Mol Cell* 2018, **71**:117–128. e113.
  43. Duda H, Arter M, Gloggnitzer J, Teloni F, Wild P, Blanco MG, Altmeyer M, Matos J: **A mechanism for controlled breakage of under-replicated chromosomes during mitosis.** *Dev Cell* 2016, **39**:740–755.
  44. Rainey MD, Bennett D, O'Dea R, Zanchetta ME, Voisin M, Seoighe C, Santocanale C: **ATR restrains DNA synthesis and mitotic catastrophe in response to CDC7 inhibition.** *Cell Rep* 2020, **32**.
  45. Teloni F, Michelena J, Lezaja A, Kilic S, Ambrosi C, Menon S, Dobrovolska A, Imhof R, Jancsak P, Baubec T, et al.: **Efficient pre-mRNA cleavage prevents replication-stress-associated genome instability.** *Mol Cell* 2019, **73**:670–683. e612.
  46. Minocherhomji S, Ying SM, Bjerregaard VA, Bursomanno S, Aeliunaite A, Wu W, Mankouri HW, Shen HH, Liu Y, Hickson ID: **Replication stress activates DNA repair synthesis in mitosis.** *Nature* 2015, **528**:286–290.
  47. Garribba L, Wu W, Ozer O, Bhowmick R, Hickson ID, Liu Y: **Inducing and detecting mitotic DNA synthesis at difficult-to-replicate loci. Mechanisms of DNA Recombination and genome rearrangements: Intersection between homologous recombination.** *DNA Replication and DNA Repair* 2018, **601**:45–58.
  48. Wu W, Bhowmick R, Vogel I, Özer Ö, Ghisays F, Thakur RS, de Leon ES, Richter PH, Ren LQ, Petrini JH, et al.: **RTEL1 suppresses G-quadruplex-associated R-loops at difficult-to-replicate loci in the human genome.** *Nat Struct Mol Biol* 2020, **27**:424–437.
  49. Ozer O, Bhowmick R, Liu Y, Hickson ID: **Human cancer cells utilize mitotic DNA synthesis to resist replication stress at telomeres regardless of their telomere maintenance mechanism.** *Oncotarget* 2018, **9**:15836–15846.
  50. Bergoglio V, Boyer AS, Walsh E, Naim V, Legube G, Lee MY, Rey L, Rosselli F, Cazaux C, Eckert KA, et al.: **DNA synthesis by Pol eta promotes fragile site stability by preventing under-replicated DNA in mitosis.** *J Cell Biol* 2013, **201**:395–408.
  51. Yang Z, Takai KK, Lovejoy CA, de Lange T: **Break-induced replication promotes fragile telomere formation.** *Genes Dev* 2020, **34**:1392–1405.
  52. Moreno A, Carrington JT, Albergante L, Al Mamun M, Haagensen EJ, Komseli ES, Gorgoulis VG, Newman TJ, Blow JJ: **Unreplicated DNA remaining from unperturbed S phases passes through mitosis for resolution in daughter cells.** In *Proceedings of the national academy of sciences of the United States of America*, vol. 113; 2016. E5757–E5764.
  53. Wu XH: **Replication stress response links RAD52 to protecting common fragile sites.** *Cancers* 2019, **11**.
  54. Bhowmick R, Minocherhomji S, Hickson ID: **RAD52 facilitates mitotic DNA synthesis following replication stress.** *Mol Cell* 2016, **64**:1117–1126.
  55. Franchet C, Hoffmann JS: **When RAD52 allows mitosis to accept unscheduled DNA synthesis.** *Cancers* 2020:12.
  56. Sotiropoulos SK, Kamileri I, Lugli N, Evangelou K, Da-Re C, Huber F, Padayachy L, Tardy S, Nicati NL, Barriot S, et al.: **Mammalian RAD52 functions in break-induced replication repair of collapsed DNA replication forks.** *Mol Cell* 2016, **64**:1127–1134.
  57. Verma P, Dilley RL, Zhang TP, Gyparakis MT, Li YW, Greenberg RA: **RAD52 and SLX4 act nonredundantly to ensure telomere stability during alternative telomere lengthening.** *Genes Dev* 2019, **33**:221–235.
  58. Wang H, Li S, Oaks J, Ren J, Li L, Wu X: **The concerted roles of FANCM and Rad52 in the protection of common fragile sites.** *Nat Commun* 2018, **9**:2791.
  59. Deng L, Wu RA, Sonnevile R, Kochenova OV, Labib K, Pellman D, Walter JC: **Mitotic CDK promotes replisome disassembly, fork breakage, and complex DNA rearrangements.** *Mol Cell* 2019, **73**:915–929. e916.
  60. Sonnevile R, Bhowmick R, Hoffmann S, Maitland N, Hickson ID, Labib K: **TRAP drives replisome disassembly and mitotic DNA repair synthesis at sites of incomplete DNA replication.** *Elife* 2019, **8**.
  61. Priego Moreno S, Jones RM, Poovathumkadavil D, Scaramuzza S, Gambus A: **Mitotic replisome disassembly depends on TRAP ubiquitin ligase activity.** *Life Sci Alliance* 2019, **2**.
  62. Wu RA, Semlow DR, Kamimae-Lanning AN, Kochenova OV, Chistol G, Hodskinson MR, Amunugama R, Sparks JL, Wang M, Deng L, et al.: **TRAP is a master regulator of DNA interstrand crosslink repair.** *Nature* 2019 Mar, **567**:267–272.
  63. Ivanova T, Maier M, Missarova A, Ziegler-Birling C, Dam M, Gomar-Alba M, Carey LB, Mendoza M: **Budding yeast complete DNA synthesis after chromosome segregation begins.** *Nat Commun* 2020, **11**:2267.
- Using budding yeast *Saccharomyces cerevisiae* as model system, this study demonstrates DNA synthesis at difficult-to-replicate regions in anaphase, after separation of sister chromatids has started, thus challenging the paradigm that DNA replication and chromosome segregation are always temporally separated.
64. Bizard AH, Hickson ID: **Anaphase: a fortune-teller of genomic instability.** *Curr Opin Cell Biol* 2018, **52**:112–119.
  65. Chan YW, Fugger K, West SC: **Unresolved recombination intermediates lead to ultra-fine anaphase bridges, chromosome breaks and aberrations.** *Nat Cell Biol* 2018, **20**:92–103.
  66. Tiwari A, Jones OA, Chan KL: **53BP1 can limit sister-chromatid rupture and rearrangements driven by a distinct ultrafine DNA bridging-breakage process.** *Nat Commun* 2018, **9**.
  67. Cuylen-Haering S, Petrovic M, Hernandez-Armendariz A, Schneider MWG, Samwer M, Blaukopf C, Holt LJ, Gerlich DW: **Chromosome clustering by Ki-67 excludes cytoplasm during nuclear assembly.** *Nature* 2020 Nov, **587**:285–290.
- This study addresses how cells separate nuclear and cytoplasmic components at the end of mitosis and identifies a chromosome clustering mechanism, which depends on the surfactant-like protein Ki-67 and displaces cytoplasmic components from the genome. Potential implications for nuclear escape of broken DNA and immune activation.
68. Samwer M, Schneider MWG, Hoefler R, Schmalhorst PS, Jude JG, Zuber J, Gerlich DW: **DNA cross-bridging shapes a single nucleus from a set of mitotic chromosomes.** *Cell* 2017, **170**:956–972. e923.
  69. Liu S, Kwon M, Mannino M, Yang N, Renda F, Khodjakov A, Pellman D: **Nuclear envelope assembly defects link mitotic errors to chromothripsis.** *Nature* 2018, **561**:551–555.
- This study reveals that spindle microtubules block assembly of nuclear envelope proteins, including nuclear pore complex components, on

lagging chromosomes in mitosis, resulting in improperly enveloped micronuclei, with implications for genome rearrangements and cytosolic sensing of self-DNA.

70. Harding SM, Benci JL, Irianto J, Discher DE, Minn AJ, Greenberg RA: **Mitotic progression following DNA damage enables pattern recognition within micronuclei.** *Nature* 2017, **548**:466–470.
  71. Mackenzie KJ, Carroll P, Martin CA, Murina O, Fluteau A, Impson DJS, Olova N, Sutcliffe H, Rainger JK, Leitch A, et al.: **cGAS surveillance of micronuclei links genome instability to innate immunity.** *Nature* 2017, **548**:461–470.
  72. Chen YA, Shen YL, Hsia HY, Tiang YP, Sung TL, Chen LY: **Extrachromosomal telomere repeat DNA is linked to ALT development via cGAS-STING DNA sensing pathway.** *Nat Struct Mol Biol* 2017, **24**:1124–1131.
  73. Nassour J, Radford R, Correia A, Fuste JM, Schoell B, Jauch A, Shaw RJ, Karlseder J: **Autophagic cell death restricts chromosomal instability during replicative crisis.** *Nature* 2019, **565**:659–663.
- This study demonstrates in the context of replicative crisis that telomere damage generates cytoplasmic DNA species, which activate the cGAS-STING pathway and trigger cGAS-STING-mediated autophagy and cell death, representing a novel tumor suppressive mechanism.
74. Maciejowski J, Chatziplis A, Dananberg A, Chu K, Toufekhtchan E, Klimczak LJ, Gordenin DA, Campbell PJ, de Lange T: **APOBEC3-dependent kataegis and TREX1-driven chromothripsis during telomere crisis.** *Nat Genet* 2020, **52**:884–890.
- Extending previous findings by the Campbell and de Lange labs that had established an important role for the cytoplasmic exonuclease TREX1 to promote resolution of dicentric chromosomes during telomere crisis, this study reveals how TREX1, in combination with cytosine deamination by APOBEC3B, promotes chromothripsis and kataegis.
75. Reislander T, Lombardi EP, Groelly FJ, Miar A, Porru M, Di Vito S, Wright B, Lockstone H, Biroccio A, Harris A, et al.: **BRCA2 abrogation triggers innate immune responses potentiated by treatment with PARP inhibitors.** *Nat Commun* 2019, **10**.
  76. Heijink AM, Talens F, Jae LT, van Gijn SE, Fehrmann RSN, Brummelkamp TR, van Vugt M: **BRCA2 deficiency instigates cGAS-mediated inflammatory signaling and confers sensitivity to tumor necrosis factor- $\alpha$ -mediated cytotoxicity.** *Nat Commun* 2019, **10**:100.
  77. Ding L, Kim HJ, Wang Q, Kearns M, Jiang T, Ohlson CE, Li BB, Xie S, Liu JF, Stover EH, et al.: **PARP inhibition elicits STING-dependent antitumor immunity in brca1-deficient ovarian cancer.** *Cell Rep* 2018, **25**:2972–2980. e2975.
  78. Chabanon RM, Muirhead G, Krastev DB, Adam J, Morel D, Garrido M, Lamb A, Henon C, Dorvault N, Rouanne M, et al.: **PARP inhibition enhances tumor cell-intrinsic immunity in ERCC1-deficient non-small cell lung cancer.** *J Clin Invest* 2019, **129**:1211–1228.
  79. Shen J, Zhao W, Ju Z, Wang L, Peng Y, Labrie M, Yap TA, Mills GB, Peng G: **PARPi triggers the STING-dependent immune response and enhances the therapeutic efficacy of immune checkpoint blockade independent of BRCAness.** *Cancer Res* 2019, **79**:311–319.
  80. Gatti M, Imhof R, Huang Q, Baudis M, Altmeyer M: **The ubiquitin ligase TRIP12 limits PARP1 trapping and constrains PARP inhibitor efficiency.** *Cell Rep* 2020, **32**:107985.
  81. Feng X, Tubbs A, Zhang C, Tang M, Sridharan S, Wang C, Jiang D, Su D, Zhang H, Chen Z, et al.: **ATR inhibition potentiates ionizing radiation-induced interferon response via cytosolic nucleic acid-sensing pathways.** *EMBO J* 2020, **39**, e104036.
  82. Reislander T, Groelly FJ, Tarsounas M: **DNA damage and cancer immunotherapy: a STING in the tale.** *Mol Cell* 2020 Oct 1, **80**:21–28.
  83. Ochs F, Karemore G, Miron E, Brown J, Sedlackova H, Rask MB, Lampe M, Buckle V, Schermelleh L, Lukas J, et al.: **Stabilization of chromatin topology safeguards genome integrity.** *Nature* 2019, **574**:571–574.
- This study uses 3D-SIM and STED microscopy to define 53BP1 nano- and higher-order microdomains around ionizing radiation-induced DNA break sites and describes a role for 53BP1 in stabilizing the topology of break-flanking chromatin.
84. Kilic S, Lezaja A, Gatti M, Bianco E, Michelena J, Imhof R, Altmeyer M: **Phase separation of 53BP1 determines liquid-like behavior of DNA repair compartments.** *EMBO J* 2019, **38**, e101379.
- This study, together with the study by Pessina et al. demonstrates that the DNA repair protein 53BP1 forms condensates around DNA lesions that show several features of phase separated compartments. Kilic et al. further demonstrate that this depends on the 53BP1 oligomerization domain and is linked to activation of the p53-p21 pathway.
85. Pessina F, Giavazzi F, Yin YD, Gioia U, Vitelli V, Galbiati A, Barozzi S, Garre M, Oldani A, Flaus A, et al.: **Functional transcription promoters at DNA double-strand breaks mediate RNA-driven phase separation of damage-response factors.** *Nat Cell Biol* 2019, **21**:1286–1299.
- This study, together with the study by Kilic et al. demonstrates that the DNA repair protein 53BP1 forms condensates around DNA lesions that show several features of phase separated compartments. Pessina et al. also provide additional evidence for a role of damage-induced long non-coding RNAs in promoting 53BP1 assembly.
86. Rai AK, Chen JX, Selbach M, Pelkmans L: **Kinase-controlled phase transition of membraneless organelles in mitosis.** *Nature* 2018, **559**:211–216.
  87. Orthwein A, Fradet-Turcotte A, Noordermeer S, Canny M, Brun C, Strecker J, Escribano-Diaz C, Durocher D: **Mitotic inhibition of the DNA double-strand break response preserves genomic integrity.** *Mol Biol Cell* 2014, **25**.
  88. Arora M, Moser J, Phadke H, Basha AA, Spencer SL: **Endogenous replication stress in mother cells leads to quiescence of daughter cells.** *Cell Rep* 2017, **19**:1351–1364.
  89. Barr AR, Cooper S, Heldt FS, Butera F, Stoy H, Mansfeld J, Novak B, Bakal C: **DNA damage during S-phase mediates the proliferation-quiescence decision in the subsequent G1 via p21 expression.** *Nat Commun* 2017 Mar 20, **8**:14728.
  90. Lezaja A, Altmeyer M: **Inherited DNA lesions determine G1 duration in the next cell cycle.** *Cell Cycle* 2018, **17**:24–32.
  91. Aymard F, Aguirrebengoa M, Guillou E, Javierre BM, Bugler B, Arnould C, Rocher V, Iacovoni JS, Biernacka A, Skrzypczak M, et al.: **Genome-wide mapping of long-range contacts unveils clustering of DNA double-strand breaks at damaged active genes.** *Nat Struct Mol Biol* 2017, **24**:353–361.
  92. Oshidari R, Huang R, Medghalchi M, Tse EYW, Ashgriz N, Lee HO, Wyatt H, Mekhail K: **DNA repair by Rad52 liquid droplets.** *Nat Commun* 2020, **11**.
- This study shows in *Saccharomyces cerevisiae* that the repair protein Rad52 forms DNA damaged-induced liquid droplets, which fuse into a larger repair center with the help of dynamic nuclear microtubule filaments. Microtubule dynamics relocate clustered lesions in liquid droplets to the nuclear periphery for repair.
93. Spies J, Lukas C, Somyajit K, Rask MB, Lukas J, Neelsen KJ: **53BP1 nuclear bodies enforce replication timing at under-replicated DNA to limit heritable DNA damage.** *Nat Cell Biol* 2019, **21**:487–497.
- This study describes a role for 53BP1 at inherited replication stress-associated DNA lesions to enforce late replication timing of under-replicated DNA and promote recombination-based, RAD52-mediated gap filling repair.
94. Schmid JA, Berti M, Walser F, Raso MC, Schmid F, Krietsch J, Stoy H, Zwicky K, Ursich S, Freire R, et al.: **Histone ubiquitination by the DNA damage response is required for efficient DNA replication in unperturbed S phase.** *Mol Cell* 2018, **71**:897–910.
- This study provides evidence for a protective role of 53BP1 and several of its upstream regulators at double-stranded DNA ends of reversed replication forks in S-phase.
95. Min J, Wright WE, Shay JW: **Clustered telomeres in phase-separated nuclear condensates engage mitotic DNA synthesis through BLM and RAD52.** *Genes Dev* 2019, **33**:814–827.
- This study, together with the study by Zhang et al. provides evidence that assembly into nuclear liquid condensates, such as ALT-associated

PML bodies (APBs), is important for several ALT features, including telomere clustering and MiDAS.

96. Zhang H, Zhao R, Tones J, Liu M, Dilley R, Chenoweth DM, Greenberg RA, Lampson MA: **Nuclear body phase separation drives telomere clustering in ALT cancer cells.** *Mol Biol Cell* 2020, mbcE19100589.

This study, together with the study by Min et al., provides evidence that assembly into nuclear liquid condensates, such as ALT-associated PML bodies (APBs), is important for several ALT features, including telomere clustering and MiDAS.

97. Vancurova M, Hanzlikova H, Knoblochova L, Kosla J, Majera D, Mistrik M, Burdova K, Hodny Z, Bartek J: **PML nuclear bodies are recruited to persistent DNA damage lesions in an RNF168-53BP1 dependent manner and contribute to DNA repair.** *DNA Repair* 2019, **78**:114–127.

98. Umbreit NT, Zhang CZ, Lynch LD, Blaine LJ, Cheng AM, Tourdot R, Sun LL, Almubarak HF, Judge K, Mitchell TJ, et al.: **Mechanisms generating cancer genome complexity from a single cell division error.** *Science* 2020, **368**:282–295.

By combining live cell microscopy and single-cell genomics, this study establishes how chromosome bridge formation during cell division errors causes aberrant replication in the next cell cycle and initiates a mutational cascade that promotes and amplifies chromothripsis.

99. Aitken SJ, Anderson CJ, Connor F, Pich O, Sundaram V, Feig C, Rayner TF, Lukk M, Aitken S, Luft J, et al.: **Pervasive lesion segregation shapes cancer genome evolution.** *Nature* 2020, **583**:265–270.

Using strand-specific tracing of mutations, this study provides evidence that mutagenic lesions are often not resolved within a single cell cycle, but rather get unevenly segregated for multiple cell generations to shape cancer genome evolution.

100. Xing M, Zhang F, Liao H, Chen S, Che L, Wang X, Bao Z, Ji F, Chen G, Zhang H, et al.: **Replication stress induces ATR/CHK1-dependent nonrandom segregation of damaged chromosomes.** *Mol Cell* 2020, **78**:714–724. e715.

This study provides evidence for nonrandom segregation of replication stress-associated genomic lesions to daughter cells during mitotic cell division.

101. Zhao PA, Sasaki T, Gilbert DM: **High-resolution Repli-Seq defines the temporal choreography of initiation, elongation and termination of replication in mammalian cells.** *Genome Biol* 2020, **21**:76.

This study, complementing the studies by Macheret et al. and Zhao et al. uses high resolution Repli-Seq to define the replication timing program in different cell types, including the mapping of termination sites and common fragile sites.

102. Ji F, Liao H, Pan S, Ouyang L, Jia F, Fu Z, Zhang F, Geng X, Wang X, Li T, et al.: **Genome-wide high-resolution mapping of mitotic DNA synthesis sites and common fragile sites by direct sequencing.** *Cell Res* 2020.

This study, together with the study by Macheret et al. maps genomic sites of MiDAS through direct sequencing, thereby globally identifying and characterizing common fragile sites and their mode of replication.

103. Macheret M, Bhowmick R, Sobkowiak K, Padayachy L, Mailler J, Hickson ID, Halazonetis TD: **High-resolution mapping of mitotic DNA synthesis regions and common fragile sites in the human genome through direct sequencing.** *Cell Res* 2020.

This study, together with the study by Ji et al. maps genomic sites of MiDAS through direct sequencing, thereby globally identifying and characterizing common fragile sites and their mode of replication.